

Effects of several factors on fungal spore germination in solid state fermentation of coprah cake

S.G. MARAKIS¹, M. LAMBRAKI¹, I. PERRAUD-GAIME², L. HANNIBAL²
AND S. ROUSSOS²

¹ Institute of General Botany, University of Athens, 157 81 Athens, Greece.

² Laboratoire de Biotechnologie, Centre ORSTOM, 911 Av. Agropolis, B.P. 5045, 34032 Montpellier cedex 1, France.

SUMMARY

Optimization of the culture conditions of *Aspergillus carbonarius* (Asca), *Aspergillus* sp (C11B52) and *Penicillium* sp (V26A25), cultured in coprah cake solid state fermentation (SSF), was undertaken, using experimental matrices. The effects of seven culture parameters (humidity = F1, initial $(\text{NH}_4)_2\text{SO}_4$, CaCl_2 and $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ concentrations = F2, F6, and F7, respectively, initial pH = F3, inoculum size = F4 and temperature = F5) on spore germination time and respiratory activities of the strains were studied. In general, the shortest lag phase was observed under the lowest values of the F1, F2, F3 and F7 parameters. Under these conditions *Aspergillus carbonarius* (Asca) and *Aspergillus* sp (C11B52) showed the highest respiratory activities (6.3% and 10.2%, respectively), while the highest respiratory activity ($\approx 5\%$) of *Penicillium* sp was observed at the lower values of F2, F3, F4 and F5 parameters.

Keywords: *Aspergillus carbonarius*, *Aspergillus* sp., *Penicillium* sp., coprah cake, spore germination, solid state fermentation, Lag phase, germination time, respiratory activity, protein enrichment.

RESUME

Effets de différents facteurs sur la germination de spores de champignons filamenteux cultivés sur tourteau de coprah en milieu solide.

MARAKIS S., LAMBRAKI M., PERRAUD-GAIME I., HANNIBAL L. ET ROUSSOS S.

L'optimisation des conditions de culture d'*Aspergillus carbonarius* (Asca), *Aspergillus* sp (C11B52) et *Penicillium* sp (V26A25), cultivés sur tourteau de coprah en milieu solide, a été réalisée en utilisant des plans d'expérience. Les effets de sept paramètres de culture (humidité = F1, concentrations initiales de $(\text{NH}_4)_2\text{SO}_4$, CaCl_2 et de $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ = F2, F6, et F7 respectivement, pH initial = F3, taille d'inoculum = F4 et température = F5) sur le temps de germination de spores et les activités respiratoires des souches ont été étudiés. Les temps les plus courts pour la germination de spores ont été obtenus pour les valeurs les plus faibles des facteurs F1, F2, F3 et F7. Sous ces conditions expérimentales *Aspergillus carbonarius* et *Aspergillus* sp. ont montré les plus grandes activités respiratoires (6,3% et 10,2% respectivement) alors que l'activité respiratoire la plus élevée ($\approx 5\%$) observée pour *Penicillium* sp. a été obtenue avec les plus faibles valeurs des paramètres F2, F3, F4 et F5.

Mots clés: *Aspergillus carbonarius*, *Aspergillus* sp., *Penicillium* sp., Tourteau de coprah, germination de spores, Phase de latence, temps de germination, activité respiratoire, enrichissement en protéines.

INTRODUCTION

The economic and social significanees of fungi, as sources of food and biologically active metabolites, stimulated the interest in their growth physiology. One of the important variables of fungal growth is spore germination time, since it determines the duration of the lag phase, and, consequently, the duration of the fungal fermentation. A prolongeard lag phase in fungal fermentation leads to increased cost of production of microbial products. It, is, thus, essential to achieve a shorter spore

germination time, either by selecting a fungal strain of short lag phase or by regulating culture conditions, e.g., tannic acid addition (Gaitis and Marakis, 1994).

A strain of *Aspergillus carbonarius*, with high tanninolytic ability, has been isolated from mouldy carob (Marakis, 1980). Although this fungus physiology has been fairly well studied for microbial protein enrichment and metabolite (enzymes) production in submerged shaker cultures (Marakis, 1980, 1985; Marakis and Diamandoglou, 1990), no study on the growth of this strain of *A. carbonarius* in SSF has been carried out, so far. A comparative study of this strain with *Aspergillus* sp and *Penicillium* sp, previously studied for protein enrichment (Roussos *et al.*, 1994) and enzyme production (Roussos *et al.*, 1995) in SSF system, is attempted.

This paper describes the effects of seven factors on the spore germination time and respiratory activities of *A. carbonarius*, *Aspergillus* sp and *Penicillium* sp, cultured in coprah cake SSF.

MATERIALS AND METHODS

MICROORGANISMS

Aspergillus carbonarius (Bainier) Thom, was isolated from mouldy carob beans (Marakis, 1980), while *Aspergillus* sp (C11B25) and *Penicillium* sp (V26A25) were obtained from the microbial collection of the Biotechnology laboratory of ORSTOM (Roussos *et al.*, 1995).

MEDIA

A mineral solution of 0.36% Na_2HPO_4 , 0.3% KH_2PO_4 and 1% urea, on initial substrate dry matter (IDM), was added to coprah residue, while $(\text{NH}_4)_2\text{SO}_4$, CaCl_2 and $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ solution of specific concentration, as shown in Table 1, were then added to obtain the desired initial humidity and mineral contents.

Table 1. Relation between code numbers (limits of factor values) and real values of the factors examined.

Factors	Limits of factor values	
	-1	1
F1 = Initial humidity (% IDM)	60	70
F2 = $(\text{NH}_4)_2\text{SO}_4$ (%IDM)	5	10
F3 = Initial pH	6	7
F4 = Inoculum size (spores/g IDM)	2×10^6	2×10^7
F5 = Temperature ($^{\circ}\text{C}$)	25	30
F6 = CaCl_2 (% IDM)	0,3	0,6
F7 = $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ (% IDM)	0,3	0,6

CULTURE CONDITIONS

The values of the seven factors studied are presented in Table 1. The optimization of the culture conditions (Table 2) was approached by using experimental matrices (De Meo *et al.*, 1985). Solid state fermentation was carried out according to the procedure of Raimbault and Alazard (1980).

SPORE GERMINATION DETERMINATION

Spores were considered to be germinated when their germ tube lengths were one and a half times the size of the spore diam (Frossard and Oertli, 1982).

MEASUREMENT OF CO_2

Carbon dioxide in the dry output air flow was continuously measured by using a gas chromatograph, as per the method of Saucedo-Castañeda *et al.* (1993).

PROTEIN EVALUATION

Proteins were estimated by total nitrogen titration, as per the total carbonization method (CHN), using a LECO SP 428 apparatus (USA), or by Kjeldhal using Kjeltabs CTQ catalyso, Thomson-Copper Ltd (Prolabo) France. The quantity of ammonium nitrogen was determined colorimetrically using the Alliance Evolution II automatic apparatus (France) and the value was deducted from the total nitrogen. Results obtained were multiplied by a factor of 5.3 for the unfermented coprah cake and by 6.25 for the fermented products (Rham, 1982).

Table 2. Experimental matrix with code numbers of the factors examined.

Expt.	Limits						
	F1	F2	F3	F4	F5	F6	F7
1	-1	-1	-1	1	1	1	-1
2	1	-1	-1	-1	-1	1	1
3	-1	1	-1	-1	1	-1	1
4	1	1	-1	1	-1	-1	-1
5	-1	-1	1	1	-1	-1	1
6	1	-1	1	-1	1	-1	-1
7	-1	1	1	-1	-1	1	-1
8	1	1	1	1	1	1	1

RESULTS AND DISCUSSION

Spore germination time and respiratory activities were found to depend on fungal strain and culture conditions. All the strain developed well enough on coprah cake in SSF, though with different respirometric profiles. This was not only because of their physiological differences (Roussos *et al.*, 1994), but also due to the culture conditions.

A. carbonarius showed the shortest lag phase (2.5 h) and the maximum respiratory activity (6.3% CO₂) after 18 h incubation in experiment No. 1, while the longest

(15 h) lag phase and the lower respiratory activity (3.75% CO₂) was observed in experiment No. 7 (Table 3). Roussos *et al* (1994) also reported a lag phase of 3 h and a respiratory activity of 7%, for the same strain and substrate. On the basis of the above data, and those presented in Fig. 1, the optimization of *A.carbonarius* growth could be achieved if a) substrate initial humidity, inoculum size and incubation temperature values are about 70%, 2×10^7 and 30°C, respectively, b) nitrogen source and pH are less than 5% and 6, respectively and c) concentrations of CaCl₂ and MgSO₄.7H₂O are lower than 0.3%. This microorganism has shown similar nutritional or physical requirements, in shaker submerged culture system (Marakis and Diamantoglou, 1990) as well as in solid state fermentation of carob pods (Lambraki *et al*, 1994), thereby indicating that this strain requires same culture conditions, independent of the cultivation system.

Table 3. Growth parameters of *A. carbonarius*, *Aspergillus* sp. and *Penicillium* sp. uner different culture conditions

Microorganisms	Expt No	Shortest germination Time (h)	Maximum CO ₂ production (%)	CO ₂ production rate (h ⁻¹)	Optimum temperature (°C)
<i>A. carbonarius</i>	1	2,5	6,3	0,37	30
<i>Aspergillus</i> sp.	1	15	10,2	0,34	30
<i>Penicillium</i> sp.	1,8	10	?	0,14	25
	2,5	?	4,8	0,14	?

Aspergillus sp. (C11B52) showed growth requirements, similar to *A. carbonarius* (Fig. 1), but the shortest lag phase (15 h) of this strain was 6 fold longer than that of *A. carbonarius*, as observed in experiment No. 1, while the longest lag period was 38 h in experiment No. 7. A temperature of 30°C was found to be optimum for both of the *Aspergillus* strains (Table 3). This result is similar to that reported by Roussos *et al* (1995) in their studies on isolation of fungal populations at different temperatures. They found that *Aspergilli* species were predominant at 30°C (= 75% of the total population), while the population of *Aspergillus* species was 25%, when the isolation temperature was 25°C. On the other hand, the maximum CO₂ production (%) in case of *Aspergillus* sp., was 1.7 fold higher than that of *A. carbonarius* (Table 3), though both the strains showed similar CO₂ production rates (0.34 h⁻¹ and 0.37 h⁻¹ for *Aspergillus* sp. and *A. carbonarius*, respectively).




Fig. 1. Protein content, CO₂ production (ml/g IDW), and lag phases of *A. carbonarius*, *Aspergillus sp.* and *Penicillium sp.* as affected by several culture parameters.

As the extent of CO₂ production indirectly reflects biomass development, the above data indicate that *Aspergillus* sp. C11B52 is most suited for biomass production as well as protein enrichment of agro-industrial byproducts/residues/wastes, while *A. carbonarius* can be of economical benefits in the production of fungal metabolites, such as enzymes, due to its shorter lag phase. Optimization of cultural conditions for each of these two strains for the above exploitation in most efficient manner could be carried out by studying other physico-chemical parameters or by further studying the presently optimized factors at closer range.

Penicillium sp. showed an intermediate (between the other two strains studied) germination time of 10-18 h, but, a comparatively low (4.8%) CO₂ production (Fig. 1). Obviously, this strain is not suitable for fermentation of the above types. Nevertheless, it is interesting to notice that the best and worst growth results for this microorganism were also recorded at experiments Nos. 1 and 7, respectively.

The fact that *Aspergillus* sp. and *Penicillium* sp. were the strains which showed maximum and minimum CO₂ production (%), respectively, is contrary to the report of Roussos *et al* (1994), who cultivated these microorganisms in the same medium. It appears that mineral concentration and aeration flow rate account for these difference in these two kinds of experiments.

Protein values were calculated from the elementary nitrogen composition, from which the amount of ammonium nitrogen was subtracted. Titration for the different samples, after 45 h of SSF, revealed that *Aspergillus* sp. cultures had a higher protein content. This was expected as this microorganism presented the highest CO₂ production (Angokai, 1993). *Penicillium* sp. also showed a high protein content, but is characterized by long phase and for this reason it is not economically beneficial.

CONCLUSIONS

Spore germination time and respiratory activity depend on the strain and culture conditions. The culture conditions used in experiment No. 1 proved the most beneficial for the growth parameters of all the fungal strains studied in coprah fermentation. Among these, *Aspergillus* sp. C11B52 is a high-biomass producer, while *A. carbonarius* completes a very quick metabolic cycle.

ACKNOWLEDGMENTS

This work was financially supported by French Government MRT/CIRAD Project N° 92 L 0401. The authors thank Dr. B.K. Lonsane, CFTRI, Mysore, India, for critical and constructive discussions.

REFERENCES

- Angokai, M.A. 1993 Enrichissement en protéines du tourteau de coprah par fermentation en milieu solide. *Mémoire DESS de Nutrition*, Université Montpellier II, France, 43p.
- De Meo, M., Laget, M., Phan-Tan-Luu, R., Mathieu, D. and Dumenil, G. 1985. Application des plans d'expériences à l'optimisation des milieux et des conditions de culture en fermentation. *Bio-Sciences* 4: 17-20.
- Frossard, R., Oertli, J.J. 1982. Growth and germination of fungal spores in guttaton fluids of Barley grown with different nitrogen sources. *Trans. Br. Mycol. Soc.* 78: 239-245.
- Gaitis, F. and Marakis, S. 1994. Tannin acid effects on spore germination time and mycelial morphology of *Aspergillus carbonarius*. *Micol. Neotrop. Appl.* 7: 5-16.
- Lambraki, M., Marakis, S. and Roussos, S. 1994. Effect of temperature and aeration flow on carob tannin degradation by *Aspergillus carbonarius* in Solid State Fermentation system. *Micol. Neotrop. Appl.* 7: 23-34.
- Marakis, S. 1980. New fungal strains for microbial protein production from carob beans. *Ph. D. Thesis*. University of Athens, Greece, p. 235.
- Marakis, S., 1985. Screening tannin-utilizing filamentous fungi for protein production from aqueous carob extract. *Cryptogamie Mycol.* 6:293-308.
- Marakis, S. and Diamandoglou, S. 1990. Fungi isolation from leaves of some Mediterranean evergreen sclerophyllous shrubs. Enzymatic activity of the isolated fungi. *Crypt. Mycol.* 11: 243-254.
- Raimbault, M. and Alazard, D. 1980. Culture method to study fungal growth in solid state fermentation. *European Appl. Microbiol. Biotechnol.* 9: 199-209.
- Rham, O. 1982. La proportion d'azote dans les protéines et le facteur de calcul protéine/azote. *Lebensm. Wiss. Technol.* 15: 226-231.

- Roussos, S., Aquiahuatl, M.D., Trejo-Hernandez, M.R., Gaimé-Perraud, I., Favela, E., Ramakrishna, M., Raimbault, M. and Viniegra-Gonzalez, G. 1995. Biotechnological management of coffee pulp - Isolation, screening, characterization, selection of caffeine-degrading fungi and natural microflora present in coffee pulp and husk. *Appl. Microbiol. Biotechnol.* 42: 756-762.
- Roussos, S., Hannibal, L., Durand, A., Diez, M., Saucedo-Castañeda, G., Montet, D. and Graille, J. 1994. Enrichissement en protéines du tourteau de coprah: sélection de champignons filamenteux en FMS. *Oléagineux*, 4: 235-247.
- Saucedo-Castañeda, G., Trejo-Hernandez M., Lonsane, B.K., Navarro, J.M., Roussos, S., Dufour, D., and Raimbault, M. 1994. On-line automated monitoring and control systems for CO₂ and O₂ in aerobic and anaerobic solid state fermentation. *Process Biochem.* 29: 13-24.